



Chem 10113, Exam 3


November 28, 2018

Name: _____
(Please Print)

1. (8 points) Write *complete Lewis electron dot formulas* for each of the following ions.



2. The simple organic compound, $\text{C}_3\text{H}_2\text{O}$, known as propynal, is very unstable but has been detected in interstellar space. Propynal has a skeletal framework as indicated in the figure below. The numbers on the figure are just labels to distinguish the carbon atoms in the following questions.



(a) (1 point) The *total* number of *valence* electrons in this molecule is _____.

(b) (2 points) In the space above, complete the Lewis electron dot formula for $\text{C}_3\text{H}_2\text{O}$.

(c) (2 points) What is the hybridization at each of the atoms?






O _____ C¹ _____ C² _____ C³ _____

(d) (1 point) The H—C¹—C² bond angle is _____ degrees.


(e) (1 point) The C²—C³—H bond angle is _____ degrees.

(f) (7 points) **Describe the bonding** in propynal, $\text{C}_3\text{H}_2\text{O}$, using **Valence Bond Theory** (i.e., hybrid atomic orbitals, etc.). **Draw and clearly label one or more pictures** to show the *types of orbitals* that you are using to form the σ and/or π bonds. Also, clearly show the 3-D structure of the molecule, including the relative orientation of the C-H, C-C-C, and C-O linkages, etc.

3. (6 points) Consider the following organic liquids. Use the letters, **A - E** to fill in the blanks in the statements below. (The same letter may be used more than once.)




- (a) _____ has the greatest surface tension.
 (b) _____ should be the least soluble in water.
 (c) _____ has the lowest vapor pressure at room temperature.
 (d) _____ has only London (dispersion) forces.
 (e) In compounds _____ and _____, the *predominant* (i.e., strongest) intermolecular interactions are *dipole-dipole forces*.
4. (3 points) Arrange the following in order of decreasing atomic radius.



5. (6 points) The following questions refer to the phase diagram of elemental iodine (I_2) as shown below (not drawn to scale).

- (a) At 200 °C and 0.20 atm is I_2 a solid, liquid, or a gas? _____
 (b) The triple point of I_2 is at $P =$ _____ atm and $T =$ _____ °C.
 (c) Which letter on the diagram indicates the critical point of I_2 ? _____
 (d) At 380 torr, I_2 boils at _____ °C.
 (e) Which letter on the diagram best represents a sublimation point? _____



6. (2 points) For a certain substance, a plot of $\ln P$ (natural log of vapor pressure, in atm) vs $1/T$ (1 over temperature, in K) affords a straight line with slope = -3675 K.
 For this substance, $\Delta H_{\text{vaporization}} =$ _____ kJ/mole.
7. (2 points) Write the *short-hand* electron configuration for rhodium (Rh).
8. (3 points) Give the *orbital diagram* for the *valence shell* electron configuration of Rh^{3+} .
9. (3 points) Circle any of the following molecules that are *non-polar*.
 PF_3 XeF_4 BrF_5 BF_3 SF_2 AsF_5
10. (3 points) Using Lewis dot symbols, illustrate the reaction of potassium and sulfur atoms to form a stable *ionic* compound.

11. (11 points) Use **Molecular Orbital Theory** as it applies to simple diatomic molecules and ions to answer the following questions.

(a) Circle any of the following molecules that are paramagnetic.



(b) The molecule NO has _____ unpaired electron(s) in the _____ energy level.

(c) The bond order of the anion NO⁻ is _____.

(d) Which has the shortest bond distance? (circle one) O₂ O₂⁺ O₂⁻ O₂²⁻

(e) Sketch the **shapes** of each of the following molecular orbitals.



12. (9 points) **SHOW ALL WORK.** The heat of combustion of propane, C₃H₈ (molar mass = 44.1), is -2220 kJ/mole. Determine the mass (in grams) of propane that is required to provide enough energy to convert 2.50 kg of ice at 273 K to water vapor at 373 K.

(Note: For H₂O, $\Delta H^\circ_{\text{fusion}} = 6.02$ kJ/mole and $\Delta H^\circ_{\text{vaporization}} = 40.7$ kJ/mole)

13. (8 points) **SHOW ALL WORK.** Write a **balanced chemical equation** for the **formation reaction** of glycine (an amino acid, structure below), and then determine the **standard heat of formation** (ΔH°_f) of glycine from the bond energy data given below.

Bond Energy (kJ/mole)

H-H	436
N-H	389
O-H	464
C-H	414
C-C	347
C-N	305
C-O	360
C=O	736
N≡N	946
O=O	498

