
## Chemistry 10123, **Exam 3** March 25, 2020

Name: \_\_\_\_\_

(Please Print)

In a detailed kinetic study of the following gas-phase reaction, [CINO<sub>2</sub>] was monitored as a function of time. The experimental data is tabulated and plotted below. Use this information to answer the following questions.
 In [CINO2]



(a) (7 points) Write the *rate law* for this reaction. In 25 words or less, explain how you arrived at your answer. (*No calculations are needed here!*)

(b) (3 points) The half-life of this reaction is \_\_\_\_\_\_ sec. (Again, no calculations!)

- (c) (3 points) SHOW ALL WORK. Determine the rate constant (k) of this reaction including proper units.
- (d) (7 points) **SHOW ALL WORK.** Determine the molar concentration of ClNO<sub>2</sub> after the reaction has been in progress for 10.0 min from the start.

2. (10 points) SHOW ALL WORK. The energy of activation of a certain reaction is 145 kJ/mole. Determine the temperature (in °C) at which the rate of this reaction is 1,000 times faster than it is at 25 °C (assuming the same initial concentrations).

 (7 points) Sketch and *correctly label* a "reaction coordinate diagram" for an <u>exo</u>thermic reaction. Such diagrams are used in the \_\_\_\_\_\_ theory of reaction rates.

4. The experimental rate law for the following reaction is found to be: rate =  $k[N_2O_5]$ .

## $N_2O_5 + O_3 \longrightarrow 2NO_2 + 2O_2$

- (a) (3 points) In 25 words or less, explain why this reaction cannot occur by a one-step mechanism.
- (b) (7 points) *Propose* a reasonable *two-step mechanism* for this reaction that is consistent with the experimental rate law. Clearly indicate which step in your mechanism is the rate-determining step. If your proposed mechanism contains an intermediate, circle it.

5. A kinetic study of the following gas-phase reaction gave the concentration vs initial rate data summarized below.

$NO + SO_3 \longrightarrow NO_2 + SO_2$				
Expt	[NO]	[SO3]	<b>initial rate</b> (mole/L·sec)	
(1)	0.150	0.250	1.65 x 10 <sup>-4</sup>	
(2)	0.450	0.450	5.35 x 10 <sup>-4</sup>	
(3)	0.150	1.350	4.81 x 10 <sup>-3</sup>	
(4)	0.150	0.450	5.35 x 10-4	

(a) (10 points) Determine the rate law for this reaction. Clearly SHOW how you arrive at your answer.

(b) (4 points) Determine the value for the **rate constant** (k) for the above reaction. **SHOW ALL WORK (including proper units)**.

6. (8 points) **SHOW ALL WORK.** For a certain reaction,  $\Delta H^{\circ} = -95.2$  kJ and  $\Delta S^{\circ} = -157$  J/K. Determine  $\Delta S_{surr}$  and  $\Delta S_{univ}$  (in J/K) for this reaction at 850 K. *Based on these results*, is the reaction spontaneous at this temperature? *Explain briefly* (but do not exceed the allotted space!).

7. (3 points) For a chemical system in a state of dynamic equilibrium,  $\Delta H$  is equal to which of the following? Circle any that apply.

 $\Delta S/R$   $E_{cell}$  log Q  $\Delta E/n$   $\Delta G$   $T\Delta S$  - RTlnK

8. (8 points) **SHOW ALL WORK.** An 80.0-L volume of an ideal gas in a cylinder with a piston is at a pressure of 3.0 atm. While the system is held at constant temperature, enough weight is placed on the piston to increase the external pressure to 10.0 atm. For the resulting process, determine q (heat) and w (work) in units of kJ, and with proper signs. (*Note*:  $1 \text{ L} \cdot \text{atm} = 101.3 \text{ J}$ )

9. In a potential hydrogen-based economy, one possible source of hydrogen would be the oxidation of natural gas with steam at high-temperature as in the following reaction. Some relevant thermodynamic data is also tabulated below.

$CH_{4(g)} + 2H_2O_{(g)} \implies CO_{2(g)} + 4H_{2(g)}$				
Compound	$\Delta H^{\circ}f(kJ/mole)$	S° (J/mole·K)		
CH <sub>4(g)</sub>	- 75	186		
$H_2O_{(g)}$	- 242	189		
CO <sub>2(g)</sub>	- 394	214		
H <sub>2(g)</sub>	0	131		

(a) (10 points) **SHOW ALL WORK.** Determine whether this reaction is spontaneous at 25 °C by calculating the appropriate thermodynamic quantity.

(b) (10 points) **SHOW ALL WORK.** Calculate the temperature (in °C) at which the equilibrium constant (K) for this reaction is equal to 100.